

AquaVac RELERA

Introduction



Company name: [Intervet / Schering-Plough Animal Health](#)
Address: **Walton Manor**
Walton
Milton Keynes
MK7 7AJ
Telephone: **01908 685685 (Customer Support Centre)**
Telephone: **01908 665050 (Switchboard)**
Fax: **01908 685555**
Email: support.uk@intervet.com
Website: www.intervet.co.uk

Presentation

The vaccine contains:

Formaldehyde-inactivated cells of *Yersinia ruckeri* (Hagerman type 1 strain) 5×10^9 cfu/mL (pre-inactivation) giving a RPS* \geq 75% ; and

Formaldehyde-inactivated cells of *Yersinia ruckeri* (SP/07/04 strain) 5×10^9 cfu/mL (pre-inactivation) giving a RPS* \geq 60% after vaccination.

*RPS: Relative Percentage Survival in Rainbow Trout.

Uses

For Rainbow Trout (*Oncorhynchus mykiss*) of 2 grams weight or over. Active immunization against Enteric Redmouth disease to reduce mortality caused by the Hagerman Type I strain and EX5 biotype of *Yersinia ruckeri*. The onset of immunity is 336 degree days (28 days at 12°C) in 5g fish. Full efficacy of the vaccine is unproven. The duration of immunity is unknown.

Dosage and administration

Immersion - Vaccination Method

1. Shake bottle well before use.
2. Dilute 1 litre of vaccine with 9 litres of clean hatchery water.
3. Drain and weigh a netful of fish and dip fish in the diluted vaccine for 30 seconds ensuring that fish are totally immersed in the vaccine.
4. After 30 seconds exposure lift net, allow to drain and return fish to holding tank.
5. Repeat until 100 kg of fish have been immersed using 1 litre of undiluted vaccine.
6. Spent vaccine is discarded after use.
7. When convenient, more than 1 litre of vaccine may be diluted in a single bath. In such cases the weight of fish which may be dipped will be 100 kg x the number of litres of vaccine used.

Contra-indications, warnings, etc

Only vaccinate healthy fish. Do not vaccinate fish suspected of having Enteric Redmouth disease.

Avoid stress at the time of the handling of fish.

Do not administer to fish intended for broodstock or broodstock.

No information is available on the interactions between AquaVacTM Relera and any other vaccine. It is therefore recommended that no other vaccine should be administered within 168 degree days (14 days at 12°C) before and after vaccination with AquaVacTM Relera. Do not mix with any other vaccine of immunological product

Withdrawal Period

Zero days

Pharmaceutical precautions

On opening the container, dilute the contents immediately and do not store.

Do not mix with any other product..

Waste vaccine and vaccine containers should be disposed of in accordance with local requirements.

On opening the container, dilute the contents immediately, and do not store. Store at 2°C to 8°C in the dark. Do not freeze.

Keep out of reach and sight of children. For animal treatment only. Dispose of waste vaccine and vaccine containers in accordance with local requirements

Legal category

POM-V

Packaging Quantities

1000ml

Further information

Nil.

Marketing authorisation number

Vm 00201/4224.

Significant Changes

New Data Sheet